

Aristichthys nobilis

Bighead carp

Advertisemen
t

Aristichthys nobilis (Richardson, 1845)

| | | |
|-------------------------|---|--|
| Family: | Cyprinidae (Minnows or carps) |  picture (Arnob_u3.jpg) by Khoo, W.L. |
| Order: | Cypriniformes (carps) | |
| Class: | Actinopterygii (ray-finned fishes) | |
| FishBase name: | Bighead carp | |
| Max. size: | 112 cm TL (male/unsexed; Ref. 40637); max. published weight: 21.3 kg (Ref. 40637) | Point map |
| Environment : | benthopelagic; potamodromous (Ref. 51243); freshwater; depth range - 5 m | |
| Climate: | temperate; 4 – 26°C; 64°N - 18°S | |
| Importance: | fisheries: highly commercial; aquaculture: commercial; aquarium: public aquariums | |
| Resilience: | Medium, minimum population doubling time 1.4 - 4.4 years(K=0.15-0.27; tm=3-5) | |
| Distribution: | Asia: China. Introduced to numerous countries and has achieved a near global distribution. However, its breeding requirements are very specialized and stocks are maintained by artificial reproduction or continuous importation. Several countries report adverse ecological impact after introduction. | |
| Gazetteer | | |
| Morphology: | Dorsal spines (total): 3 - 3; Dorsal soft rays (total): 7; Anal spines: 1 – 3; Anal soft rays: 12 – 14. Body with numerous scattered small black blotches. Keels extend from pelvic base to anus. Barbels absent. Posterior margin of last simple dorsal ray not serrated. Branched anal rays 13-14.5 (Ref. 13274). | |
| Biology: | Occurs in rivers and lakes. Feeds mainly on zooplankton. Bottom feeding fish (Ref. 6459). Marketed fresh and frozen. | |
| Red List Status: | Not in IUCN Red List (Ref. 53964) | |
| Dangerous: | potential pest | |
| Coordinator: | | |
| Main Ref: | Masuda, H., K. Amaoka, C. Araga, T. Uyeno and T. Yoshino. 1984. (Ref. 559) | |